

1. A football player is moving southward and suddenly turns eastward with the same speed to avoid an opponent. The force that acts on the player while turning is: **(2023)**  
(a) along south-west  
(b) along eastward  
(c) along northward  
(d) along north-east
2. A bullet is fired from a gun at the speed of 280 m/s in the direction  $30^\circ$  above the horizontal. The maximum height attained by the bullet is **(2023)**  
( $g = 9.8 \text{ ms}^{-2}$ ,  $\sin 30^\circ = 0.5$ )  
(a) 3000 m (b) 2800 m  
(c) 2000 m (d) 1000 m
3. A ball is projected with a velocity,  $10 \text{ ms}^{-1}$ , at an angle of  $60^\circ$  with the vertical direction, Its speed at the highest point of its trajectory will be: **(2022)**  
(a)  $5\sqrt{3} \text{ ms}^{-1}$  (b)  $5 \text{ ms}^{-1}$   
(c)  $10 \text{ ms}^{-1}$  (d) Zero
4. A particle moving in a circle of radius R with a uniform speed takes a time T to complete one revolution. If this particle were projected with the same speed at an angle  $\theta$  to the horizontal, the maximum height attained by it equals 4R. The angle of projection,  $\theta$  is then given by: **(2021)**  
(a)  $\theta = \cos^{-1} \left( \frac{\pi^2 R}{gT^2} \right)^{\frac{1}{2}}$  (b)  $\theta = \sin^{-1} \left( \frac{\pi^2 R}{gT^2} \right)^{\frac{1}{2}}$   
(c)  $\theta = \sin^{-1} \left( \frac{2gT^2}{\pi^2 R} \right)^{\frac{1}{2}}$  (d)  $\theta = \cos^{-1} \left( \frac{gT^2}{\pi^2 R} \right)^{\frac{1}{2}}$
5. A car starts from rest and accelerates at  $5 \text{ m/s}^2$ . At  $t = 4 \text{ s}$ , a ball is dropped out of a window by a person sitting in the car. What is the velocity and acceleration of the ball at  $t = 6 \text{ s}$ ? **(2021)**  
(a) 20 m/s, 0  
(b)  $20\sqrt{2} \text{ m/s}$ , 0  
(c)  $20\sqrt{2} \text{ m/s}$ ,  $10 \text{ m/s}^2$   
(d) 20 m/s,  $5 \text{ m/s}^2$
6. A ball is thrown vertically downward with a velocity of 20 m/s from the top of a tower. It hits the ground after some time with a velocity of 80 m/s. The height of the tower is: ( $g = 10 \text{ m/s}^2$ ) **(2020)**  
(a) 340 m (b) 320 m  
(c) 300 m (d) 360 m
7. A person sitting in the ground floor of a building notices through the window, of height 1.5 m, a ball dropped from the roof of the building crosses the window in 0.1 s. What is the velocity of the ball when it is at the topmost point of the window? ( $g = 10 \text{ m/s}^2$ ) **(2020 Covid Re-NEET)**  
(a) 14.5 m/s (b) 4.5 m/s  
(c) 20 m/s (d) 15.5 m/s
8. The speed of a swimmer in still water is 20 m/s. The speed of river water is 10 m/s and is flowing due east. If he is standing on the south bank and wishes to cross the river along the shortest path, the angle at which he should make his strokes w.r.t. north is given by: **(2019)**  
(a)  $30^\circ$  west (b)  $0^\circ$   
(c)  $60^\circ$  west (d)  $45^\circ$  west
9. Two particles A and B are moving in uniform circular motion in concentric circles of radii  $r_A$  and  $r_B$  with speed  $v_A$  and  $v_B$  respectively. Their time period of rotation is the same. The ratio of angular speed of A to that of B will be: **(2019)**  
(a)  $r_A : r_B$  (b)  $v_A : v_B$   
(c)  $r_B : r_A$  (d) 1 : 1
10. When an object is shot from the bottom of a long smooth inclined plane kept at an angle  $60^\circ$  with horizontal, it can travel a distance  $x_1$  along the plane. But when the inclination is decreased to  $30^\circ$  and the same object is shot with the same velocity, it can travel  $x_2$  distance. Then  $x_1 : x_2$  will be: **(2019)**  
(a)  $1 : \sqrt{2}$  (b)  $\sqrt{2} : 1$   
(c)  $1 : \sqrt{3}$  (d)  $1 : 2\sqrt{3}$

11. The angle between  $\vec{A} - \vec{B}$  and  $\vec{A} \times \vec{B}$  is ( $\vec{A} \neq \vec{B}$ ): **(2017-Gujarat)**

- (a)  $0^\circ$  (b)  $90^\circ$   
(c)  $120^\circ$  (d)  $45^\circ$

12. A ball of mass 1 kg is thrown vertically upwards and returns to the ground after 3 seconds. Another ball, thrown at  $60^\circ$  with vertical also stays in air for the same time before it touches the ground. The ratio of the two heights are: **(2017-Gujarat)**

- (a) 1 : 3 (b) 1 : 2  
(c) 1 : 1 (d) 2 : 1

13. If the magnitude of sum of two vectors is equal to the magnitude of difference of the two vectors, the angle between these vectors is: **(2016 - I)**

- (a)  $1^\circ$  (b)  $90^\circ$   
(c)  $45^\circ$  (d)  $180^\circ$

14. A particle moves so that its position vector is given by  $\vec{r} = \cos \omega t \hat{x} + \sin \omega t \hat{y}$  where  $\omega$  is a constant. Which of the following is true? **(2016 - I)**

- (a) Velocity and acceleration both are perpendicular to  $\vec{r}$   
(b) Velocity and acceleration both are parallel to  $\vec{r}$   
(c) Velocity is perpendicular to  $\vec{r}$  and acceleration is directed towards the origin  
(d) Velocity is perpendicular to  $\vec{r}$  and acceleration directed away from the origin

15. A ship A is moving Westwards with a speed of 10 km/h and a ship B 100 km South of A, is moving Northwards with a speed of 10 km/h. The time after which the distance between them becomes shortest, is: **(2015)**

- (a) 5 h (b)  $5\sqrt{2}$  h  
(c)  $10\sqrt{2}$  h (d) 0 h

16. The position vector of a particle R as a function of time is given by:

$$\vec{R} = 4 \sin(2\pi t) \hat{i} + 4 \cos(2\pi t) \hat{j}$$

Where R is in metres, t is in seconds and  $\hat{i}$  and  $\hat{j}$  denote unit vectors along x and y-direction, respectively. Which one of the following statements is wrong for the motion of particle? **(2015)**

- (a) Path of the particle is a circle of radius 4 metre  
(b) Acceleration vector is along  $-\vec{R}$   
(c) Magnitude of acceleration vector is  $\frac{v^2}{R}$  where v is the velocity of particle  
(d) Magnitude of the velocity of particle is  $8\pi$  metre/second

17. A particle is moving such that its position coordinates (x, y) are:

- (2 m, 3 m) at time t = 0,  
(6 m, 7 m) at time t = 2 s and  
(13 m, 14 m) at time t = 5 s


Average velocity vector ( $\vec{V}_{av}$ ) from t = 0 to t = 5 s is: **(2014)**

- (a)  $\frac{1}{5}(13\hat{i} + 14\hat{j})$  (b)  $\frac{7}{3}(\hat{i} + \hat{j})$   
(c)  $2(\hat{i} + \hat{j})$  (d)  $\frac{11}{5}(\hat{i} + \hat{j})$

18. A projectile is fired from the surface of the earth with a velocity of  $5 \text{ ms}^{-1}$  and angle  $\theta$  with the horizontal. Another projectile fired from another planet with a velocity of  $3 \text{ ms}^{-1}$  at the same angle follows a trajectory which is identical with the trajectory of the projectile fired from the earth. The value of the acceleration due to gravity on the planet is (in  $\text{ms}^{-2}$ ) is: (given  $g = 9.8 \text{ ms}^{-2}$ ) **(2014)**

- (a) 3.5 (b) 5.9  
(c) 16.3 (d) 110.8

19. The velocity of a projectile at the initial point A is  $(2\hat{i} + 3\hat{j}) \text{ m/s}$ . Its velocity (in m/s) at point B is: **(2013)**



- (a)  $2\hat{i} + 3\hat{j}$  (b)  $-2\hat{i} - 3\hat{j}$   
(c)  $-2\hat{i} + 3\hat{j}$  (d)  $2\hat{i} - 3\hat{j}$