
NATIONAL ELIGIBILITY CUM ENTRANCE TEST NEET (UG), 2016

Phase-2 (CODE:BB-QQ-XX)

A given nitrogen-containing aromatic compound A reacts with Sn/HCl, followed by HNO₂ to give an unstable compound B. B, on treatment with phenol, forms a beautiful coloured compound C with the molecular formula C₁₂H₁₀N₂O. The structure of compound A is



2. Consider the reaction $\label{eq:CH3CH2CH2CH2CN+NaBr} \begin{picture}(100,0) \put(0,0){\line(0,0){100}} \put(0,0){\line(0$


This reaction will be the fastest in

(1) water Polau.


methanol

N, N'-dimethylformamide (DMF) X

3. The correct structure of the product A formed in the reaction



4. Which among the given molecules ca exhibit tautomerism?



- (1) Both II and III
- (2) III only

(3) Both I and III

- (4) Both I and II
- 5. The **correct** order of strengths of the carboxylic acids

is

- (1) II > I > III
- (2) I > II > III

- (4) III > II > I
- **6.** The compound that will react most read with gaseous bromine has the formula
 - (1) C2H4
 - (2) C.H.

(4) C₄H₁₀ ⋉

- 7. Which one of the following compounds shows the presence of intramolecular hydrogen bond?
 - (1) Concentrated acetic acid

12/ H₂O₂

- (3) HCN × N-C=
- (4) Cellulose
- 8. The molar conductivity of a $0.5 \,\mathrm{mol}/\mathrm{dm}^3$ AgNO₃ with electrolytic of 5.76×10^{-3} S cm⁻¹ at of conductivity 298 K is
 - (1) $28.8 \text{ S cm}^2/\text{mol}$
 - (2) $2.88 \text{ S cm}^2/\text{mol}$

13/ 11.52 S cm²/mol

(4) $0.086 \text{ S cm}^2/\text{mol}$

- **9.)** The decomposition of phosphine (PH₃) on tungsten at low pressure is a first-order reaction. It is because the
 - (1) rate of decomposition is very slow
 - (2) rate is proportional to the surface coverage
 - (3) rate is inversely proportional to the surface coverage
 - (4) rate is independent of the surface coverage
 - 10. The coagulation values in millimoles per litre of the electrolytes used for the coagulation of As₂S₃ are given below : 1

I. (NaC1) = 52,

II. $(BaCl_2) = 0.69$, III. $(MgSO_4) = 0.22$

The correct order of their coagulating power is

- (1) III > I > II
- (2) I > II > III

(3) II > I > III (4) III > II > I,

- 11. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of 3 amperes is
 - (1) 330 minutes
 - (2) 55 minutes

(3) 110 minutes

(4) 220 minutes

- 12. How many electrons can fit in the orbital for which n = 3 and l = 1?
 - (1) 14

(3) 6

13. For a sample of perfect gas when its pressure is changed isothermally from p_i to p_f , the entropy change is given by

 $^{\circ}$ (1) $\Delta S = RT \ln \left(\frac{p_t}{p_c} \right)$

(2) $\Delta S = nR \ln \left(\frac{p_f}{p_f} \right)$

 $\Delta S = nR \ln \left(\frac{p_i}{p_f} \right)$

(4) $\Delta S = nRT \ln \left(\frac{P_f}{p_i} \right) \kappa$ = RTU VI P2

14. The van't Hoff factor (i) for a dilute aqueous solution of the strong electrolyte barium hydroxide is

M) 3

(2) 0

(3) 1

- (4) 2
- 15. The percentage of pyridine (C₅H₅N) that forms pyridinium ion (C₅H₅N⁺H) in a 0.10 M aqueous pyridine solution $C_5H_5N = 1.7 \times 10^{-9}$) is (1) 1.6% $C = \pm 1.0$ (2) 0.0060% $C = \pm 1.0$

 - JS 0.013%
 - (4) 0.77%
- 16. In calcium fluoride, having the fluorite structure, the coordination numbers for calcium ion (Ca²⁺) and fluoride ion (F⁻) are

4 and 8

- (2) 4 and 2
- (3) 6 and 6
- (4) 8 and 4
- 17. If the $E_{\rm cell}^{\circ}$ for a given reaction has a negative value, which of the following gives the **correct** relationships for the values of ΔG° and K_{eq} ?

(1) $\Delta G^{\circ} < 0$; $K_{eq} < 1$ K

- $\Delta G^{\circ} > 0$; $K_{eq} < 1$
- \sim (3) $\Delta G^{\circ} > 0$; $K_{eq} > 1$
 - (4) $\Delta G^{\circ} < 0$; $K_{eq} > 1$ ox


18. Which one of the following is incorrect for ideal solution?

$$\Delta G_{\text{mix}} = 0$$

- (2) $\Delta H_{mix} = 0$
- (3) $\Delta U_{\text{mix}} = 0$
- (4) $\Delta P = P_{\text{obs}} P_{\text{calculated by Raoult's law}} = 0$
- 19. The solubility of AgCl (s) with solubility product 1.6×10^{-10} in 0.1 M NaCl solution would be
 - (1) zero

$$1.26 \times 10^{-5} M$$

- (3) $1.6 \times 10^{-9} M$
- (4) $1.6 \times 10^{-11} M$
- 20. Suppose the elements X and Y combine to form two compounds XY_2 and X_3Y_2 . When 0.1 mole of XY_2 weighs 10 g and 0.05 mole of X₃Y₂ weighs 9 g, the atomic weights of X and Y are
 - (1) 30, 20



- 21. The number of relectrons delivered at the cathode during electrolysis by a current of 1 ampere in 60 seconds electron = 1.60×10^{-19} C)

 - (1) 7.48×10^{23} (2) 6×10^{23} (3) \times X 10

 - (3) 6×10^{20} (4) 3.75×10^{20}
- 22. Boric acid is an acid because its molecule
 - (1) combines with proton from water molecule
 - (2) contains replaceable H⁺ ion 1/4
 - (13) gives up a proton
 - (4) accepts OH from water releasing proton
- 23. AlF₃ is soluble in HF only in presence of KF. It is due to the formation of
 - (1) $K[AlF_3H]$
- (2) $K_3[AlF_3H_3]$
- $(3) K_3[AlF_6]$ (4) AlH_3

- 24. Zinc can be coated on iron to produce galvanized iron but the reverse is not possible. It is because
 - (1) zinc has higher negative electrode potential than iron
 - (2) zinc is lighter than iron
 - (3) zinc has lower melting point than iron
 - (A) zinc has lower negative electrode potential than iron
- 25. The suspension of slaked lime in water is known as
 - (1) aqueous solution of slaked lime
 - (2) limewater. Coloud, which

- (4) milk of lime
- 26. The hybridizations of atomic orbitals of nitrogen in NO2+, NO3 and NH4+ respectively
 - (1) sp^2 , sp and $sp^3 \checkmark$
 - (2) sp, sp^3 and sp^2
 - $\sqrt{21} \text{ sp}^2, \text{ sp}^3 \text{ and sp}$ sp, sp^2 and sp^3
- 27. Which of the following fluoro-compounds is most likely to behave as a Lewis base?
 - (1) SiF4
- (2) BF₃
- (4) CF₄
- 28. Which of the following pairs of ions is isoelectronic and isostructural? $(742441 \text{ Not}^{24} + 12 \text{ ClO}_3^2, \text{ SO}_3^2)$ (2) $(2) \text{ CO}_3^2$, $(2) \text{ NO}_3^2$
- (3) ClO_3^- , CO_3^{2-} (4) SO_3^{2-} , NO_3^-
- 29) In context with beryllium, which one of the following statements is incorrect?
 - (1) Its hydride is electron-deficient and polymeric.
 - (2) It is rendered passive by nitric acid.
 - (3) It forms Be 2C.
 - (4) Its salts rarely hydrolyze.

30. Hot concentrated sulphuric acid is a moderately strong oxidizing agent. Which of the following reactions does not show

oxidizing behaviour?

(A) CaF₂ + H₂SO₄ → CaSO₄ + 2HF

- (2) $Cu + 2H_2SO_4 \rightarrow CuSO_4 + SO_2 + 2H_2O_4$
- (3) $3S + 2H_2SO_4 \rightarrow 3SO_2 + 2H_2O \sim$
- (4) $C + 2H_2SO_4 \rightarrow CO_2 + 2SO_2 + 2H_2O_4$
- **31.** Which of the following pairs of d-orbitals will have electron density along the axes?
 - (1) d_{xy} , $d_{x^2-y^2}$,
 - (2) d_{z^2} , d_{xz}
 - (3) d_{xz} , d_{yz}
 - (9) d_{2} , d_{2} , d_{2}
- 32. The correct geometry and hybridization for XeF₄ are

If square planar, sp^3d^2

- (2) octahedral, sp^3d^2
- (3) trigonal bipyramidal, sp³d
 (4) planar triangle, sp³d³




- 33. Among the following, which one is a wrong statement?
 - (1) I_3^+ has bent geometry.
 - (2) PH₅ and BiCl₅ do not exist.
 - (3) $p\pi d\pi$ bonds are present in SO_2 ? SeF₄ and CH₄ have same shape.
- The **correct** increasing order of *trans-effect* of the following species is
 - (1) $CN^- > Br^- > C_6H_5^- > NH_3$
 - (2) $NH_3 > CN^- > Br^- > C_6H_5^-$
 - (3) $CN^- > C_6H_5^- > Br^- > NH_3$
 - (4) $Br^- > CN^- > NH_3 > C_6H_5^-$

Which one of the following statements related to lanthanons is incorrect?

- (1) Ce (+4) solutions are widely used as oxidizing agent in volumetric analysis.
- (2) Europium shows +2 oxidation state.
- (3) The basicity decreases as the ionic radius decreases from Pr to Lu.
- (4) All the lanthanons are much more reactive than aluminium.

- Jahn-Teller effect is not observed in high spin complexes of
 - (1) d^9
- (2) d^7
- (3) \dot{d}^{8}
- (4) d^4
- 37. Which of the following can be used as the halide component for Friedel-Crafts reaction?
 - (X) Isopropyl chloride
 - (2) Chlorobenzene X
 - (3) Bromobenzene/
 - (4) Chloroethene
- (38) In which of the following molecules, all atoms are coplanar?




- 39. Which one of the following structures represents nylon 6,6 polymer?

 $\begin{array}{c|c} & H_2 & H_{-CH_2} \\ C & C & N - CH_2 \\ H_2 & N \end{array}$

C H C H

40. In pyrrole



the electron density is maximum on

- (1) 2 and 5
- (2) 2 and 3
- (3) 3 and 4

(4) 2 and 4

- **41.** Which of the following compounds shall **not** produce propene by reaction with HBr followed by elimination or direct only elimination reaction?

 - (2) $H_2C \xrightarrow{C} CH_2$
 - (3) $H_3C C CH_2OH^{\checkmark}$

(4) H₂C=C=C

42. Which one of the following nitro-compounds **does not** react with nitrous acid?

$$(1) \quad \overset{\text{H}_3\text{C}}{\underset{\text{O}}{\bigvee}} \overset{\text{CH}_3}{\underset{\text{NO}_2}{\bigvee}}$$

- (2) $H_3C C_{NO_2}^{H_2}$
- (3) H₃C CH NO₂
- H₃C NO₂

- **43.** The central dogma of molecular genetics states that the genetic information flows from
 - (1) DNA \rightarrow RNA \rightarrow Carbohydrates
 - (2) Amino acids → Proteins → DNA
 - (3) DNA → Carbohydrates → Proteins

DNA
$$\rightarrow$$
 RNA \rightarrow Proteins

The **correct** corresponding order of names of four aldoses with configuration given below

respectively, is

- (1) D-erythrose, D-threose, L-erythrose, L-threose
- (2) L-erythrose, L-threose, L-erythrose, D-threose
- (3) D-threose, D-erythrose, L-threose, L-erythrose
- (4) L-erythrose, L-threose, D-erythrose D-threose
- 45) In the given reaction

the product P is


		0		
46.	A foreign DNA and plasmid cut by the same restriction endonuclease can be joined to form a recombinant plasmid using	(51)		primary producers of the deep-sea rothermal vent ecosystem are
			(1)	coral reefs 🗸
:-	(a) B. D.	\	(2)	green algae 🛊
	(2) Eco RI		(3)	chemosynthetic bacteria
	(3) Taq polymerase		(4)	blue-green algae
	(4) polymerase III			`
47.	Which of the following is not a component of downstream processing?	52.		ch of the following is correct for lected species?
	Expression		(1)	Small number of progeny with large size
•	(2) Separation		(2)	Large number of progeny with small size
	(3) Purification		(3)	Large number of progeny with large size
	(4) Preservation		(4)	Small number of progeny with small size
_	(i) Treservation			
48.	Which of the following restriction enzymes produces blunt ends?	53.	'-' s	sign is assigned to beneficial interaction, ign to detrimental and '0' sign to neutral raction, then the population interaction
	(1) Hind III		repr	resented by '+' '-' refers to
	(2) Sal I	`	W	parasitism
	(3) Eco RV √		(2)	mutualism
49.	(4) Xho I		(3)	amensalism
			(4)	commensalism
	Which kind of therapy was given in 1990 to a four-year-old girl with adenosine deaminase (ADA) deficiency?	54.	Whi	ch of the following is correctly matched?
	(1) Radiation therapy		(1)	Stratification—Population
	(2) Gene therapy		(2)	Aerenchyma—Opuntia
`	(3) Chemotherapy		(3)	Age pyramid—Biome
	(4) Immunotherapy		(4)	Parthenium hysterophorus—Threat to biodiversity
50.	How many hot spots of biodiversity in the world have been identified till date by	55.	Red	List contains data or information on
	orman Myers?		(1)	marine vertebrates onlyK
`	(1) 43		(2)	all economically important plants
	(2) 17		(3)	plants whose products are in
	(3) 25			international trade ^

(4) 34

(4) threatened species

- **56.** Which one of the following is **wrong** for fungi?
 - (1) They are both unicellular and multicellular.
 - (2) They are eukaryotic.
 - (3) All fungi possess a purely cellulosic cell
 - (4) They are heterotrophic.
- 57. Methanogens belong to
 - (1) Slime moulds
 - (2) Eubacteria
 - , (3) Archaebacteria
 - (4) Dinoflagellates
- 58. Select the wrong statement.
 - (1) Diatoms are microscopic and float passively in water.
 - The walls of diatoms are easily destructible.
 - (3) 'Diatomaceous earth' is formed by the cell walls of diatoms.
 - (4) Diatoms are chief producers in the oceans.
- **59.** The label of a herbarium sheet **does not** carry information on
 - height of the plant
 - (2) date of collection
 - (3) name of collector
 - (4) local names -
- **60.** Conifers are adapted to tolerate extreme environmental conditions because of
 - (1) presence of vessels
 - (2) broad hardy leaves
 - (3) superficial stomata
 - (A) thick cuticle
- **61.** Which one of the following statements is wrong?
 - (1) Laminaria and Sargassum are used as food.
 - (2) Algae increase the level of dissolved oxygen in the immediate environment.
 - Algin is obtained from red algae, and carrageenan from brown algae.
 - (4) Agar-agar is obtained from Gelidium and Gracilaria.

- 62. The term 'polyadelphous' is related to
 - (1) calyx
 - (2) gynoecium
 - 131 androecium
 - (4) corolla
 - How many plants among Indigofera, Sesbania, Salvia, Allorn, Albe, mustardy groundnut, radish, gram and turnip have stamens with different lengths in their flowers?
 - (1) Six 🗸
 - (2) Three
 - (3) Four
 - (4) Five
- 64. Radial symmetry is found in the flowers of
 - (1) Cassia 🗶
 - Brassica
 - (3) Trifolium 💢 ·
 - (4) Pişum 😿
- 65. Free-central placentation is found in
 - (1) Citrus
 - (2) Dianthus
 - (3) Argemone
 - (4) Brassica
- 66. Cortex is the region found between
 - (1) endodermis and vascular bundle
 - (2) epidermis and stele
 - (3) pericycle and endodermis
 - (4) endodermis and pith
- 67. The balloon-shaped structures called tyloses
 - are linked to the ascent of sap through xylem vessels
 - (2) originate in the lumen of vessels
 - (3) characterize the sapwood
 - (A) are extensions of xylem parenchyma cells into vessels



74. A few drops of sap were collected by cutting across a plant stem by a suitable method.

The sap was tested chemically. Which one of

the following test results indicates that it is

68. A non-proteinaceous enzyme is

(1) deoxyribonuclease

(2) lysozyme



(4) birds

equivalent to

(1) megaspore

, 127 megasporangium

(3) megasporophyll

(4) megaspore mother cell

84. The ovule of an angiosperm is technically

- for
 - (1) ensuring anaerobic conditions in the culture vessel
 - (2) purification of product
 - (3) addition of preservatives to the product
 - availability of oxygen throughout the process

- 91. A molecule that can act as a genetic material must fulfill the traits given below, except
 - (1) it should provide the scope for slow changes that are required for evolution
 - (2) it should be able to express itself in the form of 'Mendelian characters'✓
 - (3) it should be able to generate its replicat
 - (4) it should be unstable structurally and chemically
- 92. DNA-dependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the
 - (1) antistrand
 - (2) template strand
 - (3) coding strand
 - (4) alpha strand
- 93. Interspecific hybridization is the mating of
 - (1) more closely related individuals within same breed for 4-6 generations
 - (2) animals within same breed without having common ancestors
 - two different related species
 - (4) superior males and females of different breeds
- **94.**) Which of the following is **correct** regarding AIDS causative agent HIV?
- (1) HIV does not escape but attacks the acquired immune response. T- Lympho
 - (2) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase.
 - (3) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase.
 - (4) HIV is unenveloped retrovirus.
 - Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids?
 - (1) Mackerel
 - (2) Mystus
 - (3) Mangur
 - (4) Mrigala

96. Match Column-II with Column-II and select the correct option using the codes given below

Column—I Column-II

- a. Citric acid
- (i) Trichoderma
- b. Cyclosporin A
- (ii) Clostridium
- c. Statins
- (iii) Aspergillus
- d. Butyric acid
- (iv) Monascus

Codes:

- а b d (ii)K (1) (iii) (i) (iv)
- (iv) (2)(iii) (i) (ii)
- (iii) . (i) (iv) (ii)
- (iv) (4) (i) (ii) (iii)
- Biochemical Oxygen Demand (BOD) may not be a good index for pollution for water bodies (1) sugar industry -> Maleush -> Hig receiving effluents from

 - (2) domestic sewage
 - (3) dairy industry
 - (4) petroleum industry
- 98. The principle of competitive exclusion was stated by
 - (1) Verhulst and Pearl
 - (2) C. Darwin
 - (J8) G. F. Gause
 - (4) MacArthur
 - Which of the following National Parks is home to the famous musk deer or hangul?
 - (1) Dachigam National Park, Jammu &
 - (2) Keibul Lamjao National Park, Manipur
 - (3) Bandhavgarh National Park, Madhya Pradesh
 - (4) Eaglenest Wildlife Sanctuary, Arunachal Pradesh
- 100. A lake which is rich in organic waste may result in
 - mortality of fish due to lack of oxygen
 - (2) increased population of aquatic organisms due to minerals &
 - (3) drying of the lake due to algal bloom
 - (4) increased population of fish due to lots of nutrients α
- 101. The highest DDT concentration in aquatic food chain shall occur in
 - (1) eel
 - (2) phytoplankton
 - seagull
 - (4) crab

- 102. Which of the following sets of diseases is 108. Oxidative phosphorylation is caused by bacteria?
 - (1) Herpes and influenzak
 - (2) Cholera and tetanus
 - (3) Typhoid and smallpox v
 - (4) Tetanus and mumps x
- 103. Match Column-I with Column-II for housefly classification and select the correct option using the codes given below

Column—I

Column—II

- a. Family
- (i) Diptera
- b. Order
- (ii) Arthropoda
- c. Class
- (iii) Muscidae
- d. Phylum .
- (iv) Insecta


Codes:

- d h а C (1) (iv) (ii) (i) (iii)
- (ii) -(i) (iv) (iii)
- (3)(iii) (iv) (i) (ii)
- (4) (iv) (iii) (ii) (i)
- 104. Choose the correct statement.
 - (1) All Pisces have gills covered by an operculum. K
 - (2) All mammals are viviparous.
 - (3) All cyclostomes do not possess jaws and paired fins.
 - (4) All reptiles have 'a three-chambered heart. 🔀
- 105. Study the four statements (A-D) given below and select the two correct ones out of them:
 - A. Definition of biological species was given by Ernst Mayr. ~
 - B. Photoperiod does not affect reproduction in plants.
 - C. Binomial nomenclature system was given by R. H. Whittake
 - D. In unicellular organisms, reproduction is synonymous with growth.

The two correct statements are

- (1) A and B
- (2) B and C
- (3) C and D
- A and D
- In male cockroaches, sperms are stored in which part of the reproductive system?
- (1) Vas deferens ✓
- (2) Seminal vesicles
- . (3) Mushroom glands
 - (4) Testes \checkmark
- 107. Smooth muscles are
 - (1) voluntary, spindle-shaped, uninucleate
 - involuntary, fusiform, non-striated
 - (3) voluntary, multinucleate, cylindrical
 - (4) involuntary, cylindrical, striated &

- - formation of ATP by energy released from electrons removed during substrate oxidation
 - (2) formation of ATP by transfer o phosphate group from а substrate
 - (3) oxidation of phosphate group in ATP
 - (4) addition of phosphate group to ATP
- 109. Which of the following is the least likely to be involved in stabilizing the three-dimensiona folding of most proteins?
 - (1) Ester bonds
 - (2) Hydrogen bonds✓
 - (3) Electrostatic interaction
 - Hydrophobic interaction
- Which of the following describes the giver 110. graph correctly?



- (1) Exothermic reaction with energy A in absence of enzyme and B in presence of enzyme
 - Endothermic reaction with energy A in presence of enzyme and B in absence of
 - Exothermic reaction with energy A in presence of enzyme and B in absence of enzyme
 - Endothermic reaction with energy A in absence of enzyme and B in presence of enzyme
- 111. When cell has stalled DNA replication fork, which checkpoint should be predominantly activated?
 - (1) Both G_2/M and M
 - (2) G1/S
 - (3) G_2/M
 - (4) M

112	their charac	teris	of meiosis in Column—I to ic features in Column—I I rect option using the codes			
	Column—I		Column—II			
a.	Pachytene	(i)	Pairing of homologous			

- chromosomes
- b. Metaphase I
- (ii) Terminalization of chiasmata
- c. Diakinesis
- (iii) Crossing-over takes place
- d. Zygotene
- (iv) Chromosomes align at equatorial plate

Codes:

ь d а C (iv) (iii) (ii) (i) ∠ (iii) (iv) (ii) (i) (iii) (iv) (ii) (4) (ii) (iv) (iii) (i)

- **113.** Which do stimulate the hormones production of pancreatic juice and bicarbonate?
 - (1) Insulin and glucagon
 - (2) Angiotensin and epinephrine
 - (3) Gastrin and insulin
 - Cholecystokinin and secretin
- 114. The partial pressure of oxygen in the alveoli of the lungs is
 - (1) less than that of carbon dioxide (
 - (2) equal to that in the blood
 - (3) more than that in the blood
 - (4) less than that in the blood

Choose the correct statement.

- (1) Receptors do not produce potentials.
- (2) Nociceptors respond to changes in pressure.
- (3) Meissner's corpuscles are thermoreceptors.
- (4) Photoreceptors in the human eye are depolarized during darkness and become hyperpolarized in response to the light stimulus.
- 116. Graves' disease is caused due to
 - hypersecretion of adrenal gland X
 - (2) hyposecretion of thyroid gland
 - (3) hypersecretion of thyroid gland &
 - (4) hyposecretion of adrenal gland


- 117. Name the ion responsible for unmasking of active sites for myosin for cross-bridge activity during muscle contraction.
 - (1) Potassium
- (2) Calcium

Magnesium

- (4) Sodium
- 118. Name the blood cells, whose reduction in number can cause clotting disorder, leading to excessive loss of blood from the body.
 - (A) Thrombocytes-
 - (2) Erythrocytes
 - (3) Leucocytes
 - (4) Neutrophils
- 119. Name a peptide hormone which acts mainly on hepatocytes, adipocytes and enhances cellular glucose uptake and utilization.
 - (1) Gastrin
- (2) Insulin Y

(3) Glucagon

- (4) Secretin
- 120. Osteoporosis, an age-related disease of skeletal system, may occur due to
 - (1) accumulation of uric acid leading to inflammation of joints
 - (2) immune disorder affecting muscular junction leading to fatigue
 - (3) high concentration of Ca⁺⁺ and Na⁺
 - (4) decreased level of estrogen
- 121. Serum differs from blood in
 - . (1) lacking antibodies
 - (2) lacking globulins
 - . (3) lacking albumins
 - (4) lacking clotting factors
- 122. Lungs do not collapse between breaths and some air always remains in the lungs which can never be expelled because
 - (1) pressure in the lungs is higher than the atmospheric pressure
 - (2) there is a negative pressure in the lungs №
 - (3) there is a negative intrapleural pressure pulling at the lung walls
 - (4) there is a positive intrapleural pressure &
- 123. The posterior pituitary gland is not a 'true' endocrine gland because
 - (1) it secretes enzymes
 - (2) it is provided with a duct
 - (3) it only stores and releases hormones
 - (4) it is under the regulation of hypothalamus






A person can see clearly objects only when they lie between 50 cm and 400 cm from his eyes. In order to increase the maximum distance of distinct vision to infinity, the type and power of the correcting has to use, will be

- (1) convex, +0.15 diopter
- (2) convex, +2.25 diopter
- Ø concave, -0.25 diopter√

(4) concave, -0.2 diopter -1 = 0.5

A linear aperture whose width is 0.02 cm is placed immediately in front of a lens of focal length 60 cm. The aperture is illuminated normally by a parallel beam of wavelength 5×10^{-5} cm. The distance of the first dark band of the diffraction pattern from the centre of the screen i



- (2) 0·10 cm
- (3) 0·25 cm
- (4) 0·20 cm
- 138. Electrons of mass m with de-Broglie wavelength λ fall on the target in an X-ray tube. The cutoff wavelength (λ_0) of the 143. The given circuit has two ideal diodes emitted X-ray is

(1)
$$\lambda_0 = \lambda$$

$$\sqrt{2} \lambda_0 = \frac{2mc\lambda^2}{h}$$

$$(3) \ \lambda_0 = \frac{2h}{mc}$$

(4)
$$\lambda_0 = \frac{2m^2c^2\lambda^3}{h^2}$$

139. Photons with energy 5 eV are incident on a cathode C in a photoelectric cell. The maximum energy of emitted photoelectrons is 2 eV. When photons of energy 6 eV are incident on C, no photoelectrons will reach the anode A, if the stopping potential of A relative to C is

- (2) + 3 V
- (3) + 4 V
- (4) -1 V

40. If an electron in a hydrogen atom jumps from the 3rd orbit to the 2nd orbit, it emits a photon of wavelength λ . When it jumps from the 4th orbit to the 3rd orbit, the corresponding wavelength of the photon will be

$$(1) \frac{20}{13}$$

(2)
$$\frac{16}{25}\lambda$$

(3)
$$\frac{9}{16}\lambda$$

(4)
$$\frac{20}{7}\lambda$$

141 The half-life of a radioactive substance is 30 minutes. The time (in minutes) taken between 40% decay and 85% decay of the same radioactive substance is


(2) 15
$$Jn = -\lambda(\Delta T)$$


(4) 45 $2 in = \frac{in 2}{32}$

42. For CE transistor amplifier, the audio signal voltage across the collector resistance of 2 k Ω is 4 V. If the current amplification factor of loo the transistor is 100 and the base resistance is $1 k\Omega$, then the input signal voltage is

- (1) 15 mV
- (2) 10 mV
- (3/ 20 mV
- (4) 30 mV


connected as shown in the figure below. The current flowing through the res will be





- (1) 3·13 A
- (3) 10·0 A

What is the output Y in the following circuit, when all the three inputs A, B, C are first 0 and then 1?




- (3) 0, 0
- (4/ 1, 0

- 145. Planck's constant (h), speed of light in (149) vacuum (c) and Newton's gravitational constant (G) are three fundamental constants. Which of the following combinations of these has the dimension of

 - (1) $\sqrt{\frac{Gc}{h^{3/2}}}$ (2) \sqrt{hG} ML
- 146. Two cars P and Q start from a point at the same time in a straight line and their positions are represented by $x_P(t) = at + bt$ and $x_O(t) = ft - t^2$. At what time do the cars have the same velocity?
 - $\sqrt{1)} \ 2 \frac{f-a}{(1+b)}$
- (3) a+f (4) a+ 2(1+b)
- 147) In the given figure, $a = 15 \text{ m/s}^2$ represents the total acceleration of a particle moving in the clockwise direction in a circle of radius R = 2.5 m at a give instant of time. The speed of the particl



- (1) 6.2 m/s
- (2) 4.5 m/s
- (3) 5.0 m/s
- (4) 5.7 m/s
- 148. A rigid ball of mass m strikes a rigid wall at راك. 60° and gets reflected without loss of speed as shown in the figure below. The value of impulse imparted by the wall on the ball



- (2) mV
- (3) 2mV

- A bullet of mass 10 m ving horizonta with a velocity of 4 00m s⁻¹ strikes a wood block of mass 2 kg which is suspended b light inextensible string of length 5 m. A result, the centre of gravity of the block found to rise a vertical distance of 10 c The speed of the bullet after it emerges a horizontally from the block will be
 - (1) $160 \text{ m s}^{-1} = \frac{1}{3}$
 - (2) 100 m s^{-1}
 - (3) 80 m s⁻¹ 12.00= V= 4xioXio_i xi $V = 120 \text{ m s}^{-1}$ $V = 1200 \quad 20 = V$
- **150.** Two identical balls A and B having velociting of 0.5 m/s and -0.3 m/s respectively colli elastically in one dimension. The velocities Band A after the collision respectively will 1
- (1) 0.3 m/s and 0.5 m/s
 - (2) -0.5 m/s and 0.3 m/s
 - (3) 0.5 m/s and -0.3 m/s
 - 9/-0.3 m/s and 0.5 m/s
- **151.** A particle moves from a point $(-2\hat{i} + 5\hat{j})$ $(4\hat{i} + 3\hat{k})$ when a force of $(4\hat{i} + 3\hat{j})$ N applied. How much work has been done I the force? ~ a
 - (1) 2 J
 - (2) 8 J
 - (3) 11 J
- **152.** Two rotating bodies A and B of masses and 2m with moments of inertia I_A an $I_B\left(I_B>I_A\right)$ have equal kinetic energy rotation. If L_A and L_B be their angula momenta respectively, then
 - (1) $L_A > L_B$
 - (2) $L_A = \frac{L_B}{2}$
 - $(3) L_A = 2L_B$
 - $L_B > L_A$

153. A solid sphere of mass m and radius R is 157. A rectangular film of liquid is extended rotating about its diameter. A solid cylinder of the same mass and same radius is also rotating about its geometrical axis with an angular speed twice that of the sphere. The ratio of their kinetic energies of rotation

($E_{\text{sphere}} / E_{\text{cylinder}}$) will be (1) 3:1


(2) 2:3


J37 1:5 (4) 1:4

154. A light rod of length l has two masses m_1 and m_2 attached to its two ends. The moment of inertia of the system about an axis perpendicular to the rod and passing

through the centre of mass is

155. Starting from the centre of the earth having radius R, the variation of g (acceleration due to gravity) is shown by





156. A satellite of mass m is orbiting the earth (of radius R) at a height h from its surface. The total energy of the satellite in terms of g_0 , the value of acceleration due to gravity at the earth's surface, is

(1) $-\frac{2mg_0R^2}{}$

 mg_0R^2

from $(4 \text{ cm} \times 2 \text{ cm})$ to $(5 \text{ cm} \times 4 \text{ cm})$. If the work done is 3×10^{-4} J, the value of the surface tension of the liquid is

(1) 8.0 N m^{-1} (2) 0.250 N m^{-1} (3) 0.125 N m^{-1} (4) 0.2 N m^{-1} (2) 0.250 N m^{-1}

158. Three liquids of densities ρ_1 , ρ_2 and ρ_3 (with $\rho_1 > \rho_2 > \rho_3$), having the same value of surface tension T, rise to the same height in three identical capillaries. The angles of aft cos o contact θ_1 , θ_2 and θ_3 obey

(1) $\pi > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2}$

(2) $\frac{\pi}{2} > \theta_1 > \theta_2 > \theta_3 \ge 0$

 $\frac{1}{2} \frac{1}{2} \frac{1}$ $\frac{\pi}{2} < \theta_1 < \theta_2 < \theta_3 < \pi$

159. Two identical bodies are made of a material for which the heat capacity increases with temperature. One of these is at 100 °C, while the other one is at 0 °C. If the two bodies are brought into contact, then, assuming no heat loss, the final common temperature is

(1) 0 °C

(2) 50 °C

more than 50 °C

(4) less than 50 °C but greater than 0 °C

160. A body cools from a temperature 3T to 2Tin 10 minutes. The room temperature is T. Assume that Newton's law of cooling is applicable. The temperature of the body at the end of next 10 minutes

- (4) $\frac{4}{9}$ T

161. One mole of an ideal monatomic gas undergoes a process described by the equation PV^3 = constant. The heat capacity of the gas during this process is

- (4) 2R

162. The temperature inside a refrigerator is t_2 °C and the room temperature is t_1 °C. The amount of heat delivered to the room for each joule of electrical energy consumed ideally will be

> t + t $(1) \frac{t_1 + t_2}{t_1 + 273}$ (2) $\frac{t_1}{t_1 - t_2}$ (3) $\frac{t_1 + 273}{t_1 - t_2}$ (4) $\frac{t_2 + 273}{t_1 - t_2}$

A given sample of an ideal gas occupies a volume V at a pressure P and absolute temperature T. The mass of each molecule of the gas is m. Which of the following gives the density of the gas?

(1) mkT(2) P/(kT)(4) P/(kTV)(3) Pm/(kT)

164. A body of mass m is attached to the lower end of a spring whose upper end is fixed. The spring has negligible mass. When the mass m is slightly pulled down and released, it oscillates with a time period of $3 ext{ s.}$ When the mass m is increased by 1 kg, the time period of oscillations becomes $5 ext{ s.}$ The value of m in kg is

The second overtone of an open organ pipe has the same frequency as the first overtone of a closed pipe L metre long. The ength of the open pipe will be \ 3 5

166. Three sound waves of equal amplitudes have frequencies (n-1), n, (n+1). They superimpose to give beats. The number of beats produced per second will be

(1) 2 (2) 1 (3) 46 Ω (4) 26 Ω (5) 167. An electric dipole is placed at an angle of 30° 171. A long wire carrying a steady curren with an electric field intensity 2×10^5 N/C. It experiences a torque equal to 4 N m. The charge on the dipole, if the dipole length is 2 cm, is


(1) $7 \mu C$

(2), 8 mC

(**/8**) 2 mC

(4) 5 mC

A parallel-plate capacitor of area A, plat separation d and capacitance C is filled wit four dielectric materials having dielectri constants k_1 , k_2 , k_3 and k_4 as shown in th figure below. If a single dielectric material; to be used to have the same capacitance Ci this capacitor, then its dielectric constant is given by




(1) $\frac{1}{k} = \frac{1}{k_1} + \frac{1}{k_2} + \frac{1}{k_3} + \frac{A}{2k}$

(2) $k = k_1 + k_2 + k_3 + 3k_4$

(3) $k = \frac{2}{3}(k_1 + k_2 + k_3) + 2k_4$

 $\int_{0}^{2} \frac{2}{k} = \frac{3}{k_{1} + k_{2} + k_{3}} + \frac{1}{k_{4}}$

169. The potential difference $(V_A - V_B)$ betwee the points A and B in the given figure is



170. A filament bulb (500 W, 100 V) is to be use in a 230 V main supply. When a resistance is connected in series, it works perfectly an the bulb consumes 500 W. The value of R is

> (1) 13Ω (2) 230 Ω

into a circular loop of one turn. The magnet field at the centre of the loop is B. It is the bent into a circular coil of n turns. Th magnetic field at the centre of this coil n turns will be

(1) $2n^2B$


- 172. A bar magnet is hung by a thin cotton thread 176. The in a uniform horizontal magnetic field and is in equilibrium state. The energy required to rotate it by 60° is W. Now the torque required to keep the magnet in this new position is
 - 2W(1)


- 173. An electron is moving in a circular path under the influence of a transverse magnetic field of 3.57×10^{-2} T. If the value of e/m is 1.76×10^{11} C/kg, the frequency of revolution of the electron is
 - (1) 6·28 MHz
- (2) 1 GHz
- (3) 100 MHz
- (4) 62·8 MHz



- Which of the following combinations should be selected for better tuning of an L-C-R circuit used for communication?
- (1) $R = 25 \Omega$ L = 1.5 H, $C = 45 \mu$ F
- (2) $R = 20 \Omega$, L = 1.5 H, $C = 35 \mu F$
- (3) $R = 25 \Omega$, L = 2.5 H, $C = 45 \mu F$
- (4) $R = 15 \Omega$ L = 3.5 H. $C = 30 \mu$ F
- 175. A uniform magnetic field is restricted within a region of radius r. The magnetic field changes with time at a rate $\frac{dB}{dt}$. Loop 1 of

radius R > r encloses the region r and loop 2 of radius R is outside the region of magnetic field as shown in the figure below. Then the e.m.f. generated is





- $-\frac{d\vec{B}}{dt}\pi r^2$ in loop 1 and zero in loop 2
- (2) zero in loop 1 and zero in loop 2
- (3) $-\frac{dB}{dt}\pi r^2$ in loop 1 and $\sqrt{}$ $-\frac{d\vec{B}}{dt}\pi r^2 \text{ in loop 2} \qquad \qquad \cancel{S} \frac{2\sqrt{n}}{n+1}$
- (4) $-\frac{d\vec{B}}{dt}\pi R^2$ in loop 1 and zero in loop 2

- differences potential across resistance, capacitance and inductance are 80 V, 40 V and 100 V respectively in an L-C-R circuit. The power factor of this circuit is
 - (1) 1.0
- PO,
- (3) 0.5
- A 100 Ω resistance and a capacitor of 100 Ω reactance are connected in series across a 220 V source. When the capacitor is 50% charged, the peak value of the displacement current is
 - (1) $11\sqrt{2}$ A
- (2) 2·2 A 💛 📆
- (3) 11 A
- (4) 4·4 A
- 178. Two identical glass ($\mu_g = 3/2$) equiconvex lenses of focal length f each are kept in contact. The space between the two lenses is filled with water ($\mu_w = 4 / 3$). The focal length of the combination is

(3) f (4)
$$\frac{4}{4}$$

- 179. An air bubble in a glass slab with refractive 🎺 index 1.5 (near normal incidence) is 5 cm deep when viewed from one surface and 3 cm deep when viewed from the opposite face. The thickness (in cm) of the slab is
 - (1) 16
- (3) 10
- (3/ 12 75 5 % 31 b)
- 180. The interference pattern is obtained with two coherent light sources of intensity ratio n. In the interference pattern, the ratio

$$I_{\frac{\max}{I_{\max}} + I_{\min}}$$

will be

- $(1) \quad \frac{2\sqrt{n}}{(n+1)^2} \quad \underline{\text{Imq}} \, \Lambda$

- $(4) \quad \frac{\sqrt{n}}{(n+1)^2}$