

1. Match List I with List II. (2024)

List I (Conversion)		List II (Number of Faraday required)	
A.	1 mole of $\text{H}_2\text{O}$ to $\text{O}_2$	I.	3F
B.	1 mol of $\text{MnO}_4^-$ to $\text{Mn}^{2+}$	II.	2F
C.	1.5 mol of Ca from molten $\text{CaCl}_2$	III.	1F
D.	1 mol of $\text{FeO}$ to $\text{Fe}_2\text{O}_3$	IV.	5F

Choose then correct answer from the options given below:

- (a) A-III, B-IV, C-I, D-II  
(b) A-II, B-III, C-I, D-IV  
(c) A-III, B-IV, C-II, D-I  
(d) A-II, B-IV, C-I, D-III
2. Mass in grams of copper deposited by passing 9.6487 A current through a voltmeter containing copper sulphate solutions for 100 seconds is:  
(Given : Molar mass of Cu :  $63 \text{ g mol}^{-1}$ .  
(1F = 96487 C) (2024)
- (a) 0.315 g  
(b) 31.5 g  
(c) 0.0315 g  
(d) 3.15 g
3. The  $E^\ominus$  values for  
 $\text{Al}^+/\text{Al} = +0.55 \text{ V}$  and  $\text{Tl}^+/\text{Tl} = -0.34 \text{ V}$   
 $\text{Al}^{3+}/\text{Al} = -1.66 \text{ V}$  and  $\text{Tl}^{3+}/\text{Tl} = +1.26 \text{ V}$   
Identify the incorrect statement (2023)
- (a) Al is more electropositive than Tl  
(b)  $\text{Tl}^{3+}$  is a good reducing agent than  $\text{Tl}^{1+}$   
(c)  $\text{Al}^+$  is unstable in solution  
(d)  $\text{Al}^+$  can be easily oxidised Tl than  $\text{Tl}^{3+}$
4. Molar conductance of an electrolyte increase with dilution according to the equation:  $\Lambda_m = \Lambda_m^\circ - A\sqrt{c}$   
Consider the following four statements:

- A. This equation applies to both strong and weak electrolytes.  
B. Value of the constant A depends upon the nature of the solvent.  
C. Value of constant A is same for both  $\text{BaCl}_2$  and  $\text{MgSO}_4$ .  
D. Value of constant A is same for both  $\text{BaCl}_2$  and  $\text{Mg}(\text{OH})_2$ .

Which of the above statements are correct? (2023)

- (a) A and B only (b) A, B and C only  
(c) B and C only (d) B and D only

5. The correct value of cell potential in volt for the reaction that occurs when the following two half cells are connected, is  
 $\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s}), E^\circ = -0.44 \text{ V}$   
 $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$   
 $E^\circ = +1.33 \text{ V}$  (2023)

- (a) +1.77 V (b) +2.65 V  
(c) +0.01 V (d) +0.89 V

6. The conductivity of centimolar solution of KCl at  $25^\circ\text{C}$  is  $0.0210 \text{ ohm}^{-1} \text{ cm}^{-1}$  and the resistance of the cell containing the solution at  $25^\circ\text{C}$  is 60 ohm. The value of cell constant is- (2023)

- (a)  $3.28 \text{ cm}^{-1}$  (b)  $1.26 \text{ cm}^{-1}$   
(c)  $3.34 \text{ cm}^{-1}$  (d)  $1.34 \text{ cm}^{-1}$

7. Given below are two statements: One is labelled as Assertion A and the other is labelled as Reason R:

**Assertion:** In equation  $\Delta_{\text{rG}} = -nFE_{\text{cell}}$ , value of  $\Delta_{\text{rG}}$  depends on n.

**Reason:**  $E_{\text{cell}}$  is an intensive property and  $\Delta_{\text{rG}}$  is an extensive property.

In the light of the above statements, choose the correct answer from the options given below: (2023)

- (a) Both A and R are true but R is NOT the correct explanation of A.  
(b) A is true but R is false.  
(c) A is false but R is true.  
(d) Both A and R are true R is the correct explanation of A.

8. Two half cell reactions are given below:  
 $\text{Co}^{3+} + e^- \rightarrow \text{Co}^{2+}$ ,  $E_{\text{Co}^{2+}/\text{Co}^{3+}}^\circ = -1.81 \text{ V}$   
 $2\text{Al}^{3+} + 6e^- \rightarrow 2\text{Al(s)}$ ,  $E_{\text{Al}/\text{Al}^{3+}}^\circ = +1.66 \text{ V}$   
 The standard EMF of a cell with feasible redox reaction will be: **(2022)**  
 (a)  $-3.47 \text{ V}$  (b)  $+7.09 \text{ V}$   
 (c)  $+0.15 \text{ V}$  (d)  $+3.47 \text{ V}$
9. Standard electrode potential for the cell with cell reaction  
 $\text{Zn(s)} + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{Cu(s)}$   
 is  $1.1 \text{ V}$ . Calculate the standard Gibbs energy change for the cell reaction.  
 (Given:  $F = 96487 \text{ C mol}^{-1}$ ) **(2022)**  
 (a)  $-200.27 \text{ J mol}^{-1}$   
 (b)  $-200.27 \text{ kJ mol}^{-1}$   
 (c)  $-212.27 \text{ kJ mol}^{-1}$   
 (d)  $-212.27 \text{ J mol}^{-1}$
10. At  $298 \text{ K}$ , the standard electrode potentials of  $\text{Cu}^{2+}/\text{Cu}$ ,  $\text{Zn}^{2+}/\text{Zn}$ ,  $\text{Fe}^{2+}/\text{Fe}$  and  $\text{Ag}^+/\text{Ag}$  are  $0.34 \text{ V}$ ,  $-0.76 \text{ V}$ ,  $-0.44 \text{ V}$  and  $0.80 \text{ V}$ , respectively.  
 On the basis of standard electrode potential, predict which of the following reaction can not occur?  
 (a)  $\text{CuSO}_4(\text{aq}) + \text{Zn(s)} \rightarrow \text{ZnSO}_4(\text{aq}) + \text{Cu(s)}$   
 (b)  $\text{CuSO}_4(\text{aq}) + \text{Fe(s)} \rightarrow \text{FeSO}_4(\text{aq}) + \text{Cu(s)}$   
 (c)  $\text{FeSO}_4(\text{aq}) + \text{Zn(s)} \rightarrow \text{ZnSO}_4(\text{aq}) + \text{Fe(s)}$   
 (d)  $2\text{CuSO}_4(\text{aq}) + 2\text{Ag(s)} \rightarrow 2\text{Cu(s)} + \text{Ag}_2\text{SO}_4(\text{aq})$
11. Given below are half cell reactions:  
 $\text{MnO}_4^- + 8\text{H}^+ + 5e^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$ ,  
 $E_{\text{Mn}^{2+}/\text{MnO}_4^-}^\circ = -1.510 \text{ V}$   
 $\frac{1}{2}\text{O}_2 + 2\text{H}^+ + 2e^- \rightarrow \text{H}_2\text{O}$   
 $E_{\text{O}_2/\text{H}_2\text{O}}^\circ = +1.223 \text{ V}$   
 will the permanganate ion,  $\text{MnO}_4^-$  liberate  $\text{O}_2$  from water in the presence of an acid? **(2022)**  
 (a) Yes, because  $E_{\text{cell}}^\circ = +0.287 \text{ V}$   
 (b) No, because  $E_{\text{cell}}^\circ = -0.287 \text{ V}$   
 (c) Yes, because  $E_{\text{cell}}^\circ = +2.733 \text{ V}$   
 (d) No, because  $E_{\text{cell}}^\circ = -2.733 \text{ V}$
12. Find the emf of the cell in which the following reaction takes place at  $298 \text{ K}$   
 $\text{Ni(s)} + 2\text{Ag}^+(0.001 \text{ M}) \rightarrow \text{Ni}^{2+}(0.001 \text{ M}) + 2\text{Ag(s)}$   
 (Given that  $E_{\text{cell}}^\circ = 1.05 \text{ V}$ ,  $\frac{2.303 RT}{F} = 0.059$  at  $298 \text{ K}$ ) **(2022)**  
 (a)  $1.0385 \text{ V}$  (b)  $1.385 \text{ V}$   
 (c)  $0.9615 \text{ V}$  (d)  $1.05 \text{ V}$
13. The molar conductance of  $\text{NaCl}$ ,  $\text{HCl}$  and  $\text{CH}_3\text{COONa}$  at infinite dilution are  $126.45$ ,  $426.16$  and  $91.0 \text{ S cm}^2 \text{ mol}^{-1}$  respectively. The molar conductance of  $\text{CH}_3\text{COOH}$  at infinite dilution is. Choose the right option for your answer. **(2021)**  
 (a)  $390.71 \text{ S cm}^2 \text{ mol}^{-1}$   
 (b)  $698.28 \text{ S cm}^2 \text{ mol}^{-1}$   
 (c)  $540.48 \text{ S cm}^2 \text{ mol}^{-1}$   
 (d)  $201.28 \text{ S cm}^2 \text{ mol}^{-1}$
14. The molar conductivity of  $0.007 \text{ M}$  acetic acid is  $20 \text{ S cm}^2 \text{ mol}^{-1}$ . What is the dissociation constant of acetic acid? Choose the correct option. **(2021)**  

$$\left[ \begin{array}{l} \Lambda_{\text{H}^+}^\circ = 350 \text{ S cm}^2 \text{ mol}^{-1} \\ \Lambda_{\text{CH}_3\text{COO}^-}^\circ = 50 \text{ S cm}^2 \text{ mol}^{-1} \end{array} \right]$$
  
 (a)  $2.50 \times 10^{-4} \text{ mol L}^{-1}$   
 (b)  $1.75 \times 10^{-5} \text{ mol L}^{-1}$   
 (c)  $2.50 \times 10^{-5} \text{ mol L}^{-1}$   
 (d)  $1.75 \times 10^{-4} \text{ mol L}^{-1}$
15. On electrolysis of dil sulphuric acid using Platinum (Pt) electrode, the product obtained at anode will be: **(2020)**  
 (a) Oxygen gas (b)  $\text{H}_2\text{S}$  gas  
 (c)  $\text{SO}_2$  gas (d) Hydrogen gas
16. The number of Faradays (F) required to produce  $20 \text{ g}$  of calcium from molten  $\text{CaCl}_2$  (Atomic mass of  $\text{Ca} = 40 \text{ g mol}^{-1}$ ) is: **(2020)**  
 (a) 2 (b) 3  
 (c) 4 (d) 1
17. Identify the reaction from following having top position in EMF series (Std. red. Potential) according to their electrode potential at  $298 \text{ K}$ . **(2020 Covid Re-NEET)**  
 (a)  $\text{Fe}^{2+} + 2e^- \rightarrow \text{Fe(s)}$   
 (b)  $\text{Au}^{3+} + 3e^- \rightarrow \text{Au(s)}$   
 (c)  $\text{K}^+ + 1e^- \rightarrow \text{K(s)}$   
 (d)  $\text{Mg}^{2+} + 2e^- \rightarrow \text{Mg(s)}$
18. In a typical fuel cell, the reactants (R) and product (P) are **(2020 Covid Re-NEET)**  
 (a)  $\text{R} = \text{H}_{2(\text{g})}$ ,  $\text{O}_{2(\text{g})}$ ;  $\text{P} = \text{H}_2\text{O(l)}$   
 (b)  $\text{R} = \text{H}_{2(\text{g})}$ ,  $\text{O}_{2(\text{g})}$ ,  $\text{Cl}_{2(\text{g})}$ ;  $\text{P} = \text{HClO}_{4(\text{aq})}$   
 (c)  $\text{R} = \text{H}_{2(\text{g})}$ ,  $\text{N}_{2(\text{g})}$ ;  $\text{P} = \text{NH}_{3(\text{aq})}$   
 (d)  $\text{R} = \text{H}_{2(\text{g})}$ ,  $\text{O}_{2(\text{g})}$ ;  $\text{P} = \text{H}_2\text{O}_{2(\text{l})}$

19. For a cell involving one electron  $E^\circ_{\text{cell}} = 0.59 \text{ V}$  at 298 K, the equilibrium constant for the cell reaction is: **(2019)**  
 [Given that  $\frac{2.303RT}{F} = 0.059 \text{ V}$  at  $T = 298 \text{ K}$ ]  
 (a)  $1.0 \times 10^2$  (b)  $1.0 \times 10^5$   
 (c)  $1.0 \times 10^{10}$  (d)  $1.0 \times 10^{30}$
20. For the cell reaction  
 $2\text{Fe}^{3+}(\text{aq}) + 2\text{I}^{-}(\text{aq}) \rightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{I}_2(\text{aq})$   
 $E^\circ_{\text{cell}} = 0.24 \text{ V}$  at 298 K. The standard Gibbs energy ( $\Delta_r G^\circ$ ) of the cell reaction is:  
 [Given that Faraday constant  $F = 96500 \text{ C mol}^{-1}$ ] **(2019)**  
 (a)  $-46.32 \text{ kJ mol}^{-1}$   
 (b)  $-23.16 \text{ kJ mol}^{-1}$   
 (c)  $46.32 \text{ kJ mol}^{-1}$   
 (d)  $23.16 \text{ kJ mol}^{-1}$
21. Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:
- $$\begin{array}{c} \text{BrO}_4^- \xrightarrow{1.82 \text{ V}} \text{BrO}_3^- \xrightarrow{1.5 \text{ V}} \text{HBrO} \\ \text{Br} \xleftarrow{1.0652 \text{ V}} \text{Br}_2 \xleftarrow{1.595 \text{ V}} \end{array}$$
- Then the species undergoing disproportionation is: **(2018)**  
 (a)  $\text{BrO}_3^-$  (b)  $\text{BrO}_4^-$   
 (c)  $\text{HBrO}$  (d)  $\text{Br}_2$
22. In the electrochemical cell  
 $\text{Zn}|\text{ZnSO}_4(0.01\text{M})||\text{CuSO}_4(1.0\text{M})|\text{Cu}$ , the emf of this Daniel cell is  $E_1$ . When the concentration of  $\text{ZnSO}_4$  is changed to 1.0 M and that of  $\text{CuSO}_4$  changed to 0.01 M, the emf changes to  $E_2$ . From the following, which one is the relationship between  $E_1$  and  $E_2$ ? (Given,  $\frac{RT}{F} = 0.059$ ) **(2017-Gujarat)**  
 (a)  $E_2 = 0 \neq E$  (b)  $E_1 = E_2$   
 (c)  $E_1 < E_2$  (d)  $E_1 > E_2$
23. Given that  
 $\Lambda_m^\circ = 133.4, 5 \text{ cm}^2 \text{ mol}^{-1} (\text{AgNO}_3)$ ;  
 $\Lambda_m^\circ = 149.9 \text{ S cm}^2 \text{ mol}^{-1} (\text{KCl})$ ;  
 $\Lambda_m^\circ = 144.9 \text{ S cm}^2 \text{ mol}^{-1} (\text{KNO}_3)$  the molar conductivity at infinite dilution for  $\text{AgCl}$  is: **(2017-Gujarat)**  
 (a)  $132 \text{ S cm}^2 \text{ mol}^{-1}$  (b)  $140 \text{ S cm}^2 \text{ mol}^{-1}$   
 (c)  $138 \text{ S cm}^2 \text{ mol}^{-1}$  (d)  $134 \text{ S cm}^2 \text{ mol}^{-1}$
24. The zinc/silver oxide cell is used in electric watches. The reaction is as following:  
 $\text{Zn}^{2+} + 2\text{e}^{-} \rightarrow \text{Zn}$ ;  $E^\circ = -0.760 \text{ V}$   
 $\text{Ag}_2\text{O} + \text{H}_2\text{O} + 2\text{e}^{-} \rightarrow 2\text{Ag} + 2\text{OH}^{-}$ ;  
 $E^\circ = 0.344 \text{ V}$   
 If  $F$  is  $96,500 \text{ C mol}^{-1}$ ,  $\Delta G^\circ$  of the cell will be: **(2017-Gujarat)**  
 (a)  $413.21 \text{ kJ mol}^{-1}$   
 (b)  $113.072 \text{ kJ mol}^{-1}$   
 (c)  $-213.072 \text{ kJ mol}^{-1}$   
 (d)  $4313.082 \text{ kJ mol}^{-1}$
25. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of 3 amperes is: **(2016-II)**  
 (a) 220 minutes (b) 330 minutes  
 (c) 55 minutes (d) 110 minutes
26. Zinc can be coated on iron to produce galvanized iron but the reverse is not possible. It is because: **(2016-II)**  
 (a) Zinc has lower negative electrode potential than iron  
 (b) Zinc has higher negative electrode potential than iron  
 (c) Zinc is lighter than iron  
 (d) Zinc has lower melting point than iron
27. The number of electrons delivered at the cathode during electrolysis by a current of 1 ampere in 60 seconds is: **(2016-II)**  
 (charge on electron =  $1.60 \times 10^{-19} \text{ C}$ )  
 (a)  $3.75 \times 10^{20}$  (b)  $7.48 \times 10^{23}$   
 (c)  $6 \times 10^{23}$  (d)  $6 \times 10^{20}$
28. If the  $E^\circ_{\text{cell}}$  for a given reaction has a negative value, which of the following gives the correct relationships for the values of  $\Delta G^\circ$  and  $K_{\text{eq}}$ ? **(2016-II)**  
 (a)  $\Delta G^\circ < 0$ ;  $K_{\text{eq}} > 1$   
 (b)  $\Delta G^\circ < 0$ ;  $K_{\text{eq}} < 1$   
 (c)  $\Delta G^\circ > 0$ ;  $K_{\text{eq}} < 1$   
 (d)  $\Delta G^\circ > 0$ ;  $K_{\text{eq}} > 1$
29. The molar conductivity of a  $0.5 \text{ mol dm}^{-3}$  solution of  $\text{AgNO}_3$  with electrolytic conductivity of  $5.76 \times 10^{-3} \text{ S cm}^{-1}$  at 298 K is: **(2016-II)**  
 (a)  $0.086 \text{ S cm}^2 \text{ mol}^{-1}$   
 (b)  $28.8 \text{ S cm}^2 \text{ mol}^{-1}$   
 (c)  $2.88 \text{ S cm}^2 \text{ mol}^{-1}$   
 (d)  $11.52 \text{ S cm}^2 \text{ mol}^{-1}$

30. The pressure of  $H_2$  required to make the potential of  $H_2$  electrode zero in pure water at 298 K is: **(2016-I)**  
 (a)  $10^{-4}$  atm (b)  $10^{-14}$  atm  
 (c)  $10^{-12}$  atm (d)  $10^{-10}$  atm
31. A device that converts energy of combustion of fuels like hydrogen and methane, directly into electrical energy is known as: **(2015)**  
 (a) Electrolytic cell (b) Dynamo  
 (c) Ni-Cd cell (d) Fuel cell
32. The pair of compounds that can exist together is: **(2014)**  
 (a)  $HgCl_2, SnCl_2$  (b)  $FeCl_2, SnCl_2$   
 (c)  $FeCl_3, KI$  (d)  $FeCl_3, SnCl_2$
33. Using the Gibbs energy change,  $\Delta G^0 = +63.3 \text{ kJ}$ , for the following reaction,  $Ag_2CO_3(s) \rightleftharpoons 2Ag^+(aq) + CO_3^{2-}(aq)$  the  $K_{sp}$  of  $Ag_2CO_3(s)$  in water at  $25^\circ\text{C}$  is: **(2014)**  
 ( $R = 8.314 \text{ JK}^{-1}\text{mol}^{-1}$ )  
 (a)  $8.0 \times 10^{-12}$  (b)  $2.9 \times 10^{-3}$   
 (c)  $7.9 \times 10^{-2}$  (d)  $3.2 \times 10^{-26}$
34. When 0.1 mol  $MnO_4^{2-}$  is oxidized, the quantity of electricity required to completely oxidise  $MnO_4^{2-}$  to  $MnO_4^-$  is: **(2014)**  
 (a) 96500 C (b)  $2 \times 96500 \text{ C}$   
 (c) 9650 C (d) 96.50 C
35. The weight of silver (atomic weight = 108) displaced by a quantity of electricity which displaces 5600 mL of  $O_2$  at STP will be: **(2014)**  
 (a) 10.8 g (b) 54.0 g
- (c) 108.0 g (d) 5.4 g
36. At  $25^\circ\text{C}$ , molar conductance of 0.1 molar aqueous solution of ammonium hydroxide is  $9.54 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  and at infinite dilution its molar conductance is  $238 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ . The degree of ionization of ammonium hydroxide at the same concentration and temperature is: **(2013)**  
 (a) 2.080% (b) 20.800%  
 (c) 4.008% (d) 40.800%
37. A hydrogen gas electrode is made by dipping platinum wire in a solution of HCl of pH = 10 and by passing hydrogen gas around the platinum wire at one atm pressure. The oxidation potential of electrode would be? **(2013)**  
 (a) 0.059 V (b) 0.59 V  
 (c) 0.118 V (d) 1.18 V
38. A button cell used in watches functions as following  
 $Zn(s) + Ag_2O(s) + H_2O(l) \rightleftharpoons 2Ag(s) + Zn^{2+}(aq) + 2OH^-(aq)$ .  
 If half cell potentials are  
 $Zn^{2+}(aq) + 2e^- \rightarrow Zn(s)$  ;  $E^0 = -0.76 \text{ V}$   
 $Ag_2O(s) + H_2O(l) + 2e^- \rightarrow 2Ag(s) + 2OH^-(aq)$   
 $E^0 = 0.34 \text{ V}$  . The cell potential will be: **(2013)**  
 (a) 1.10 V (b) 0.42 V  
 (c) 0.84 V (d) 1.34 V