





DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

T.B.C.: AAO-1/24

Sl. No.

14337

TEST BOOKLET

ASSISTANT AGRICULTURE OFFICER PAPER-I

K-48

Time Allowed: 2 hrs.

Test Booklet Series





INSTRUCTIONS TO CANDIDATES

- 1. IMMEDIATELY AFTER COMMENCEMENT OF THE EXAMINATION, YOU SHOULD CHECK THAT THIS TEST BOOKLET DOES NOT HAVE ANY UNPRINTED OR TORN OR MISSING PAGES OR ITEMS ETC. IF SO, GET IT REPLACED BY A COMPLETE TEST BOOKLET OF SAME SERIES ISSUED TO YOU.
- 2. ENCODE CLEARLY THE TEST BOOKLET SERIES A, B, C OR D; AS THE CASE MAY BE, IN THE APPROPRIATE PLACE IN THE ANSWER SHEET USING BALL POINT PEN (BLUE OR BLACK).
- 3. You have to enter your Roll No. on the Test Booklet in the Box provided along side. DO NOT write anything else on the Test Booklet.
- 4. YOU ARE REQUIRED TO FILL UP & DARKEN ROLL NO., TEST BOOKLET / QUESTION BOOKLET SERIES IN THE ANSWER SHEET AS WELL AS FILL UP TEST BOOKLET / QUESTION BOOKLET SERIES AND SERIAL NO. AND ANSWER SHEET SERIAL NO. IN THE ATTENDANCE SHEET CAREFULLY. WRONGLY FILLED UP ANSWER SHEETS ARE LIABLE FOR REJECTION AT THE RISK OF THE CANDIDATE.
- STORT OF THE TAX OF TH
- 5. This Test Booklet contains 100 items (questions). Each item (question) comprises four responses (answers). You have to select the correct response (answer) which you want to mark (darken) on the Answer Sheet. In case, you feel that there is more than one correct response (answer), you should mark (darken) the response (answer) which you consider the best. In any case, choose ONLY ONE response (answer) for each item (question).
- 6. You have to mark (darken) all your responses *ONLY* on the separate Answer Sheet provided, by using BALL POINT PEN (BLUE OR BLACK). See instructions in the Answer Sheet.
- 7. All items (questions) carry equal marks. All items (questions) are compulsory. Your total marks will depend only on the number of correct responses (answers) marked by you in the Answer Sheet. There shall be negative marking for wrong answers with a deduction of 0.25 marks for every wrong answer.
- 8. Before you proceed to mark (darken) in the Answer Sheet the responses to various items (questions) in the Test Booklet, you have to fill in some particulars in the Answer Sheet as per instructions sent to you with your Admission Certificate.
- 9. After you have completed filling in all your responses on the Answer Sheet and after conclusion of the examination, you should hand over to the Invigilator the *Answer Sheet* issued to you. You are allowed to take with you the candidate's copy / second page of the Answer Sheet along with the Test Booklet, after completion of the examination, for your reference.
- 10. Sheets for rough work are appended in the Test Booklet at the end.

DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO



Test Prime

ALL EXAMS, ONE SUBSCRIPTION



70,000+ Mock Tests



600+ Exam Covered



Personalised Report Card



Previous Year Papers



Unlimited Re-Attempt



500% Refund

















ATTEMPT FREE MOCK NOW





- 1. Which tillage implement is suitable for breaking the hard pan formed at plough sole layer of soil profile?
 - (A) Indigenous plough
 - (B) Chisel Plough
 - (C) MB Plough
 - (D) Blade harrow
- 2. A tillage system when 15-30% soil surface remains under crop residues cover after planting is called
 - (A) Zero tillage
 - (B) Conservation tillage
 - (C) Reduced tillage
 - (D) Intensive tillage
- 3. The crop(s) known as pseudo cereals is/are
 - (A) Amaranthus cruentus
 - (B) Fagopyrum esculentum
 - (C) Chenopodium quinoa
 - (D) All of these
- 4. Which one of the following nutrient elements plays essential role in pollen tube germination?
 - (A) Nitrogen
 - (B) Sulphur
 - (C) Boron
 - (D) Molybdenum

- 5. Which one of the following nutrient elements play beneficial roles in sugarcane crop, but is not an essential nutrient element?
 - (A) Sodium
 - (B) Nickel
 - (C) Vanadium
 - (D) Silicon
- 6. Which one of the following fertilizers supplies secondary nutrients along with primary nutrient?
 - (A) Di-ammonium phosphate
 - (B) Muriate of potash
 - (C) Single super phosphate
 - (D) Nano urea
- 7. For which category of following plants, the available soil water decreases gradually from FC to PWP resulting into gradual decrease in yield?
 - (A) Xerophytes like cactus
 - (B) Shallow rooted plants like onion and garlic
 - (C) Perennial plants like trees
 - (D) Upland crops like maize in sandy loam soils
- 8. In which method of irrigation, evaporation loss of water is minimized considerably by adopting land configuration?
 - (A) Check basin
 - (B) Border strip
 - (C) Sprinkler
 - (D) Ridge and Furrow





- 9. Which one of the following instruments is useful to know the right time of applying irrigation to field crops grown in non-saline sandy loam soil?
 - (A) Tensiometer
 - (B) Piezometer
 - (C) Lux meter
 - (D) Hygrometer
- 10. Which one of the following herbicides is commonly used for pre-emergence application in upland crops?
 - (A) 2,4-D
 - (B) Isoproturon
 - (C) Glyphosate
 - (D) Pendimethalin
- 11. The ratio between total cropped area and actual net cultivated area expressed in percentage is known as
 - (A) Cropping intensity
 - (B) Harvest index
 - (C) Rotational intensity
 - (D) Land equivalent ratio
- 12. The practice(s) is/are meant for midseason correction as a contingent management to overcome moistures stress due to the dry spell in dryland agriculture include(s)
 - (A) urea top dressing
 - (B) thinning of plants
 - (C) re-sowing for gap filling
 - (D) all of these

- 13. All decisions and activities of a farming system emanate from
 - (A) Farm household
 - (B) Farm implements and machineries
 - (C) Farm livestock
 - (D) Farm enterprises
- 14. Yearly sequence and spatial arrangements of crops and crops and fallow on a given piece of land is termed as
 - (A) Cropping system
 - (B) Cropping pattern
 - (C) Cropping sequence
 - (D) Crop rotation
- 15. A rotation of arable crops requiring annual cultivation and artificial pasture occupying field for two years or longer is termed as
 - (A) Alley cropping
 - (B) Inter cropping
 - (C) Ley farming
 - (D) Crop rotation
- **16.** Which one of the following is the key underlying issue in the concept of sustainable agriculture?
 - (A) Social equity
 - (B) Gender equity
 - (C) Inter-generational equity
 - (D) Economic equity





- 17. Which one of the following sets of agricultural practices could be ideal to achieve sustainable agriculture?
 - (A) Zero tillage, crop residue retention and crop rotation
 - (B) Intensive tillage, crop residue burning, mono cropping
 - (C) Zero tillage, crop residue removal, crop rotation
 - (D) Intensive tillage, crop residue incorporation, crop rotation
- 18. Principles of health, principles of ecology, principles of fairness and principles of care as four basic principles of organic agriculture are proposed by
 - (A) APEDA
 - (B) IFOAM
 - (C) ICAR
 - (D) FAO
- 19. On which aspect, the organic certification is different from other certification?
 - (A) product certification
 - (B) accreditation
 - (C) standards
 - (D) process certification

- 20. Which one of the following accreditation programmes has been formulated by the Government of India for the certification of organic farming and has been recognized by many countries internationally?
 - (A) IFOAM
 - (B) JAS
 - (C) NPOP
 - (D) NOP
- 21. What is the duration of conversion period required for the establishment of a certified organic farm?
 - (A) 1 year
 - (B) 2-3 years
 - (C) 4-5 years
 - (D) 8-10 years
- 22. Which one of the following types of planting materials is completely prohibited in certified organic farming?
 - (A) Certified organic seed and plant materials
 - (B) Uncertified organic seed and plant materials
 - (C) Chemically untreated conventional seed and plant materials
 - (D) Genetically engineered seeds, transgenic plants or plant materials





- **23.** Which one of the following practices does not address management of soil fertility in a certified organic farming?
 - (A) Cultivation of legumes
 - (B) Use of green manures
 - (C) Cultivation of shallow rooted crops
 - (D) Crop rotation
- **24.** Which one of the following growth stages of rice is most sensitive to nitrogen deficiency?
 - (A) Panicle initiation
 - (B) Flag leaf
 - (C) Milk stage
 - (D) Flowering
- **25.** What is the seed rate of transplanted rice?
 - (A) 10 kg/ha
 - (B) 20 kg/ha
 - (C) 30 kg/ha
 - (D) 50 kg/ha
- **26.** Which one of the following growth stage of wheat is most sensitive to soil moisture and nitrogen supply?
 - (A) Crown root initiation
 - (B) Tillering
 - (C) Flowering
 - (D) Late jointing

- 27. In which one of the following cereal crops, earthing up is generally an essential practice particularly when cultivated in rainy season?
 - (A) Barley
 - (B) Wheat
 - (C) Paddy
 - (D) Maize
- 28. Shubhra and Kripa are varieties of
 - (A) Chickpea
 - (B) Wheat
 - (C) Cotton
 - (D) Lentil
- 29. Which one of the following is the most important agrotechnique for the cultivation of rabi pulse crops in India particularly under rainfed condition?
 - (A) Deep summer tillage
 - (B) Timely sowing
 - (C) Urea top dressing
 - (D) Thinning
- 30. Which combination of following fertilizers should be more useful in oilseed crops?
 - (A) Urea, DAP, MOP
 - (B) Urea, SSP, MOP
 - (C) Mixed fertilizer (10-26-26), Urea
 - (D) DAP only

K-48 [P.T.O.





- 31. "Pegging" is an important growth stage and is found in case of
 - (A) Groundnut
 - (B) Sesame
 - (C) Cotton
 - (D) Mustard
- **32.** The distance between two rows of sugarcane planted in autumn season is generally followed as
 - (A) 40 cm
 - (B) 60 cm
 - (C) 75 cm
 - (D) 90 cm
- **33.** The dormancy of seed tuber of potato can be broken with the treatment of
 - (A) Thiourea + IAA
 - (B) Thiourea + IBA
 - (C) Thiourea+GA
 - (D) Thiourea + CO,
- 34. What should be right stage of harvesting jute crop for better quality as well as higher yield of jute fibre?
 - (A) pre-flowering stage
 - (B) flowering stage
 - (C) small pod stage
 - (D) matured pod stage

- **35.** What is the seed rate of fodder maize sown by broadcast method?
 - (A) 20 kg/ha
 - (B) 40 kg/ha
 - (C) 60 kg/ha
 - (D) 80 kg/ha
- **36.** Disease free plants in micro propagation can be obtained through
 - (A) Meristem culture
 - (B) Anther culture
 - (C) Embryo culture
 - (D) Cell culture
- 37. Example of an aggregate fruits is:
 - (A) Pineapple
 - (B) Custard apple
 - (C) Apple
 - (D) Orange
- **38.** "Pusa Nanha" dwarf variety of Papaya is developed through
 - (A) Hybridization
 - (B) Mutation
 - (C) Selection
 - (D) Heterosis



39.	Which Mango variety is suitable for high density planting?	43.	Guava is generally trained in system
	(A) Sindhu		(A) modified leader
	(B) Amrapali		(B) central leader
	(C) Mallika		(C) open leader
	(D) Ambika		(D) modified central leader
40.	Cultivated strawberry is:	44.	Sapota is commercially propagated by
	(A) Diploid(B) Octaploid		(A) inarching
	(C) Hexaploid (D) Triploid		(B) cleft grafting(C) whip grafting
			(D) four-flap grafting
41.	Vegetative propagation of date palm is through (A) Runners (B) Slips (C) Suckers	45.	Among the following which is the best planting material for banana? (A) Water suckers (B) Sword suckers
	(D) Stolen		(C) Pseudo suckers(D) Peeper suckers
42.	Indicator plant for tristeza virus in citrus is :	46.	The inception of ripening in grapes is
	(A) Citrus aurantifolia(B) Citrus reticulata(C) Citrus maxima		(A) Verasion phase (B) Lag phase
	(D) Citrus medica		(C) Bloom phase(D) Skin tanning phase





4	o ·			1
47.	Onion	15	propagated	by
	~		L - L - D	- 5

- (A) bulb
- (B) seed
- (C) layering
- (D) cutting

48. Pusa Early Synthetic is a cultivar of

- (A) Cabbage
- (B) Knolkhol
- (C) Cauliflower
- (D) Turnip

49. Buck eye rot is associated with

- (A) Tomato
- (B) Cabbage
- (C) Onion
- (D) French bean

50. Kochia is a

- (A) summer season annual flower
- (B) summer season annual foliage
- (C) winter season annual flower
- (D) winter season annual foliage

51. Zoysia japonica is commonly used in

- (A) Hedge
- (B) Edge
- (C) Lawn
- (D) Pergola

52. Degreening of citrus fruit is done by the application of

- (A) Ethylene
- (B) Pectin methyl esterase
- (C) Naphthalene
- (D) Gibrallic acid

53. The physiological disorder 'Calyx splitting' is associated with

- (A) Marigold
- (B) Carnation
- (C) Rose
- (D) Lily

54. White Flowered Gourd is also known as

- (A) Bottle gourd
- (B) Ash gourd
- (C) Sponge gourd
- (D) Apple gourd



- 5. Commercially litchi is propagated through .
 - (A) Air layering
 - (B) Stool layering
 - (C) Serpentine layering
 - (D) Trench layering
- 6. 'Bordeaux mixture' was discovered by
 - (A) E.F. Smith
 - (B) Needham
 - (C) P.M.A. Millardet
 - (D) None of the above
- 7. Sesame phyllody disease is caused by
 - (A) Bacteria
 - (B) Virus
 - (C) Fungi
 - (D) Phytoplasma
- 8. The vector of Yellow vein mosaic of bhindi is
 - (A) Aphids
 - (B) Mites
 - (C) Whiteflies
 - (D) Jassids

- **59.** Bacterial genus having strong pectolytic activity is
 - (A) Erwinia sp
 - (B) Peudomonas sp
 - (C) Ralstonia sp
 - (D) Xanthomonas sp
- 60. The causal agent of Khaira disease of rice is
 - (A) Cu deficiency
 - (B) Zn deficiency
 - (C) Mn deficiency
 - (D) Na deficiency
- 61. An antifungal antibiotic is
 - (A) Streptomycin
 - (B) Kasugamycin
 - (C) Tetracycline
 - (D) None of these
- 62. Most of the plant viruses have their nucleic acid as
 - (A) ssRNA
 - (B) dsRNA
 - (C) ssDNA
 - (D) dsDNA





- 63. Red rust of mango caused by
 - (A) fungi
 - (B) bacteria
 - (C) virus
 - (D) algae
- 64. Irish Famine is associated with
 - (A) Early blight of potato
 - (B) Leaf roll of potato
 - (C) Late blight of potato
 - (D) None of these
- 65. The pathogen of white rust of mustard belongs to the order
 - (A) Uredinales
 - (B) Ustilaginales
 - (C) Peronosporales
 - (D) Erysiphales
- 66. The pathogen causing black rot of crucifers generally enters into the host through
 - (A) Root primordial
 - (B) Lenticels
 - (C) Hydathodes
 - (D) Stomata

- 67. Which option does not fall unde cultural controls of insects?
 - (A) Crop rotation
 - (B) Tillage
 - (C) Weeding
 - (D) Bagging
- **68.** Androconia found in the wings of som insects are
 - (A) Specialized scales
 - (B) Bristles
 - (C) Spots
 - (D) Specialized glands
- **69.** Which stored grain pest requires lowes moisture content for survival
 - (A) Rhizopertha dominica
 - (B) Tribolium castaneum
 - (C) Tribolium confusum
 - (D) Trogoderma granarium
- **70.** Tomato pinworm belongs to the family
 - (A) Pterophoridae
 - (B) Agromyzidae
 - (C) Pyralidae
 - (D) Gelechidae

AAO-1/24/A

1

10

K-48





- 71. Rossetting of flowers in cotton is due to
 - (A) Pink boll worm
 - (B) American Boll worm
 - (C) Spotted boll worm
 - (D) All of these
- 72. The non-hymenopteran parasitoid recommended for pest management is
 - (A) Apanteles sp.
 - (B) Brachymeria sp.
 - (C) Epipyrops sp.
 - (D) Pachycrepoideus sp
- 73. Physapodan insects are
 - (A) White fly
 - (B) Thrips
 - (C) Leafhoppers
 - (D) Aphid
- 74. From a crop damage function $Y = 729.5 4.25 x^2 + 3.5x$ then find the unit insect damage per unit area (where Y = Kg/ha; x = insect damage per plant)
 - (A) 8 kg/ha
 - (B) 7 kg/ha
 - (C) 6 kg/ha
 - (D) 5 kg/ha

- 75. Find the EIL of BPH if an average of 20 hoppers/hill caused 500 kg/ha yield loss. The cost of management and market price were Rs. 3000/ha & Rs. 1500 / Qnt resp. and efficiency of chemical treatment is 80%:
 - (A) 20 hoppers/hill
 - (B) 15 hoppers/hill
 - (C) 10 hoppers/hill
 - (D) 8 hoppers/hill
- **76.** Higher threshold level in pest management is practiced for
 - (A) Crops having high compensatory ability
 - (B) Crops having low compensatory ability
 - (C) Crops having moderate compensatory ability
 - (D) Crops having no compensatory ability
- 77. Damage boundary is the lowest level of injury which is caused?
 - (A) Economic damage
 - (B) Damage equal to EIL
 - (C) Damage equal to ETL
 - (D) Measurable damage
- 78. "Cry for help" theory in tritropic interaction shows positive interaction in between
 - (A) Host plant and Pests
 - (B) Predators and Parasites
 - (C) Both Pests and Predators
 - (D) Both Host and Predators

1





- 79. If the discharge rate of a nozzle is 30 litters /hr, swath width (with normal stretched arm) is 200 cm and speed of the operator is 3 km/hr, then required volume /ha is:
 - (A) 300 lit
 - (B) 400 lit
 - (C) 500 lit
 - (D) 600 lit
- **80.** Reducing the width of a bund is a common practice to combat which infestation in the field?
 - (A) Heiroglyphuss banian
 - (B) Bandicota bengalensis
 - Agrotis ipsilon
 - (D) Holotrichia consanguinea
- Smothering of ear head in paddy is the damage symptoms of what?
 - Leptocorisa acuta
 - Brebhenia rehi
 - Heiroglyphus banian
 - (D) Scirpophaga incertulas

- Dark spiral streak on cane rind is found t be made of a series punctures lying sid by side like beads in a rosary symptom i caused by:
 - Chilo infuscatellus (A)
 - Chilo sacchariphagus indicus (B)
 - (C) Scirpophaga nivella
 - (D) Acigona steniellus
- The giant cell formation in infected plant 83. roots is due to
 - Meloidogyne sp (A)
 - (B) Pythium sp
 - (C) Anguina sp
 - (D) Virus
- Heterodera avenue is a
 - (A) Root knot nematode
 - Cyst nematode
 - (C) Lesion nematode
 - (D) Lance nematode

12 K-48 AAO-1/24/A



5.	'Onion Bloat' disease is caused by ——— (A) Ditylenchus dipsaci (B) Tylenchulus semipenetrans (C) Rotylenchulus reniformis (D) Pratylanchus sp.	89. Undegummed silk thread spun by the silkworm; it consists of two filaments that are cemented together by silk sericin is called as (A) Bave (B) Brin (C) Fibril (D) Cuite
6.	Non-domesticated species of silk worm that produce Chinese tasar is: (A) Antheraea mylitta (B) Antheraea royeli (C) Antheraea proyeli (D) Antheraea pernyi	90. 'Black spots' on the lateral body aspect of the infected silkworm larvae is due to infection of (A) Virus (B) Bacteria (C) Fungus
 8. 	The weight of cocoons required to produce a kilogram of raw silk is called (A) Renditta (B) Denier (C) Pongee (D) Noil A silk filament of 1000 m length and weighing 28 centigram then the estimated Denier of the filament is: (A) 2.80 (B) 2.52 (C) 1.82 (D) 3.00	91. Cutting followed by planting of new saplings of tree is (A) afforestation (B) deforestation (C) reforestation (D) none of these 92. Production of woody plants combined with pasture is referred to (A) Silvipastoral system (B) Taungya system (C) Kangeyan system (D) Agrisilvipastoral system





- 93. One of the most common features noticed in forest areas is
 - (A) Ecological niche shift
 - (B) Highly productive
 - (C) Ecological succession
 - (D) Poor diversity index
- **94.** Which one of the following is not a green house gas
 - (A) Carbon dioxide
 - (B) Methane
 - (C) Nitrous oxide
 - (D) Hydrogen
- 95. Which among the following is used for artificial rain making
 - (A) Silver iodide
 - (B) Sodium chloride
 - (C) Dry ice
 - (D) All of these
- **96.** Which one of the following is a day neutral plant?
 - (A) Barley
 - (B) Cotton
 - (C) Sunflower
 - (D) Maize

- 97. Line connected between the places with equal sunshine hours is called
 - (A) Isotach
 - (B) Isohyet
 - (C) Isohel
 - (D) Isoprecet
- 98. Out of total radiation received by the earth from the Sun the waves significan to plant life are
 - (A) UV, Radio Waves and Visible
 - (B) Radio, UV and Infrared
 - (C) Visible, Radio and Infrared
 - (D) Visible, UV and Infrared
- 99. In weather satellite the sensor mainly used is
 - (A) Photography
 - (B) Line scan
 - (C) Microwave system
 - (D) Both Line scan and microwave
- 100. Doldrums are
 - (A) High pressure areas
 - (B) Low pressure areas
 - (C) Temperate areas
 - (D) All of these





SPACE FOR ROUGH WORK











